



ADVANCED  
ORTHOPAEDIC  
SPECIALISTS

## Post-Operative Instructions **Routine Elbow Arthroscopy**

### Surgical Dressing:

1. A sterile operative dressing has been applied to your elbow. If possible, we prefer that it not be altered or changed for 72 hours following surgery. If it comes loose or is soiled, then it may be changed.
2. The elbow is filled with water to perform your surgery. It is normal for the elbow to drain/ooze water for the first day or so. This is usually blood-tinged so don't be alarmed. If you have questions, please call.
3. Three days after your surgery you may shower. I ask that you cover the elbow during the shower with saran wrap or some similar material to keep the elbow dry. You will find several (amount will vary depending on the procedure performed.) small surgical incisions with sutures place regular band-aids over each incision. You may then change the band-aids daily after showering.
4. Do not soak the elbow in bathtub, hot tub, or pool until told it is ok in clinic. Do not expose the wounds to any creek, lake, river, etc. for at least 1 month.

### Diet:

As tolerated you may resume a normal diet at your own pace. We advise though that the day of surgery you avoid any heavy meals. Our recommendation is that you begin with a liquid diet the day of surgery, and advance as tolerated. It is not uncommon to have nausea after the surgery because of the pain medicine and anesthetics used during your surgery, so be careful with what you eat and go slowly.

### Sling:

We will give you instructions on your limitations and restrictions as you move along. If you have a splint preventing motion of your elbow then it is to remain in place until seen in clinic. If you do not have a splint then you may begin moving your elbow as tolerated, and I encourage you to do so to decrease stiffness, pain, and to control swelling. You may remove the sling and begin gentle range of motion exercises as soon as you're ready, we recommend the starting gently the day after surgery. You should not do any pushing, pulling, or lifting with your operative arm. Nothing heavier than a coffee cup should be held in your operative arm until we see you back in clinic unless otherwise stated. It is ok to eat using your operative arm. You should wear your sling for sleeping and when out in public for the first several days, you may then stop using it as you are able. Ideally, you should be out of the sling when we see you back for your first follow-up appointment.

### Medications:

Patients' medication requirements vary after surgery. You have given medications for pain, nausea, and constipation prevention. Take the pain medicine if you need it, if you take it, take the stool softener as instructed. Take the nausea medicine if you need it.

Numbing medicine has been placed in your elbow at the conclusion of your surgery. Additionally, a nerve block may have been performed by the anesthesiologist. It is normal to have some level of discomfort after any surgery. If you have no pain or minimal pain, then it likely means the numbing medicines are working. It is important that you take pain medicine as the numbing medicine starts to wear off, this will prevent you from having extreme pain once it's worn off completely. If you are awake and alert, take one pain pill every 4 hours until the numbing medicines have worn completely off. Do not take the medicines if you are excessively drowsy. This helps prevent the pain from "piling up" and will significantly limit the amount of pain medicine you require in the following days.

If you have any allergies, or develop any adverse reactions to any of the medicines prescribed for you DO NOT TAKE THEM. Call and adjustments can be made.

If tolerated, and approved by your medical doctor, you may take Advil, Motrin, or similar anti-inflammatories to aid in pain control. It is ok to take these in addition to the pain medicines I have prescribed. The medicine you have been prescribed for pain likely has Tylenol in it--DO NOT TAKE ADDITIONAL TYLENOL. I advise you to talk with your pharmacist regarding potential drug interactions with other medications you are taking prior to taking any new medicines including those written for you for this surgery.

### Ice:

Controlling swelling is one of the best ways to control pain. If an ice machine has been given to you use it as instructed. Remember to always keep a towel, t-shirt, or sheet between the ice pad and your skin. If this is not done, your skin can become frostbitten.

If you do not have an ice machine you may use ice packs, gel-packs, or similar cold therapy. In general, I recommend 1 hour on, and 1 hour off. Remember to keep something between the ice and your skin.

### Concerns:

It is common to have a low-grade fever for the first 24-48 hours after any surgery. If you have a fever that does not respond to Tylenol or that is >100.5 you should call. If you have any other signs of symptoms of infection you should call. While the risk of blood clots is very low following elbow surgery, we still recommend that you ambulate as normally as possible immediately. Also, we encourage you to do ankle pump exercises while in bed for the first few days following surgery. Of course, if you have any questions or concerns, I would rather you call than worry.

Follow-Up: In general, you will be seen in clinic 2 weeks following surgery. Your appointment may be scheduled prior to your surgery, if not please call to arrange it.